

Strong approximation for the supermarket model

Malwina J Luczak ^{*}& James Norris [†]

February 1, 2008

Abstract

We prove three strong approximation theorems for the “supermarket” or “join the shortest queue” model – a law of large numbers, a jump process approximation and a central limit theorem. The estimates are carried through rather explicitly. This allows us to estimate each of the infinitely many components of the process in its own scale and to exhibit a cut-off in the set of active components which grows slowly with the number of servers.

Key words and phrases. Supermarket model, join the shortest queue, law of large numbers, diffusion approximation, exponential martingale inequalities.

MSC (2000). Primary: 60K25 . Secondary: 60F15.

1 Introduction

The supermarket model is a system of N single-server queues. Customers arrive as a Poisson process of rate $N\lambda$, where $\lambda \in (0, 1)$. Each customer examines d queues, chosen randomly from all N queues, where $d \geq 2$, and joins the shortest of these d queues, choosing randomly if the shortest queue is not unique. The service times of all customers are independent rate 1

^{*}Department of Mathematics, London School of Economics, Houghton Street, London WC2A 2AE, UK. e-mail: malwina@planck.lse.ac.uk

[†]Statistical Laboratory, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WB, UK. e-mail: J.R.Norris@statslab.cam.ac.uk

exponential random variables. We will be concerned with the behaviour of this model when λ and d are fixed, over a finite time interval $[0, t_0]$, as $N \rightarrow \infty$. We shall consider the case when the system starts in some well-behaved state with low server loads (in a sense to be made precise below).

This model has attracted attention because it turns out that the choice offered to customers, even if $d = 2$, dramatically reduces queue lengths; see [5, 13, 15]; and in particular, the length of the longest queue, see [10, 11]. Given that our analysis relies on N being very much larger than d , the model does not describe well the behaviour of a real supermarket. Rather it serves as an example where a simple dynamic routing rule leads to a greatly improved performance, which is of interest in the context of communications networks.

Our results provide strong approximations for the supermarket model, and include a law of large numbers and a diffusion approximation. In arriving at these results we have developed techniques to establish weak convergence of a sequence of Markov processes X^N in infinitely many dimensions, where the jumps of X^N are of order N^{-1} and occur at a rate of order N . The classical results for fluid limits are set in a finite-dimensional context. We make essential use of the fact that the number of “active” components in X^N grows only very slowly with N . We have used direct and quantitative methods based on exponential martingales and strong approximation of Poisson processes by Brownian motion. These methods seem well-suited to deal with such “almost finite-dimensional” Markov processes. Earlier results for this model include laws of large numbers in [5, 6, 13, 15], quantitative concentration of measure estimates in [11] and a central limit theorem [7]. See also [9] for a preliminary version of the law of large numbers presented in the present paper (for a more general range of initial conditions).

The limiting behaviour of the supermarket model as $N \rightarrow \infty$ may conveniently be described in terms of the vector $X_t = (X_t^k : k \in \mathbb{N})$, where X_t^k denotes the proportion of all N queues having at least k customers at time t . The process $X = (X_t)_{t \geq 0}$ is a Markov chain. We will suppose throughout that $X_0 = x_0$ with x_0 non-random and we will suppress the dependence of X on N to lighten the notation. Now X has the form of a density dependent Markov chain such as considered by Ethier and Kurtz in [4], Chapter 11. Thus one might expect to be able to find a deterministic process $(x_t)_{t \geq 0}$ and a Gaussian process $(\gamma_t)_{t \geq 0}$ such that

$$X_t = x_t + O(N^{-1/2}), \quad X_t = x_t + N^{-1/2}\gamma_t + O(\log N/N).$$

However the number of non-zero components in X grows with N so the

standard theory does not apply.

We will see that, for small initial data, the component X^k has a scale $a_k = \lambda^{1+d+\dots+d^{k-1}}$, which of course decays very rapidly with k . Thus the number of queues having at least k customers is of order Na_k . We can find m of order $\log \log N$ such that Na_m is of order 1. Thus we can exhibit a cut-off in the number of active components which grows only slowly with N . Below the cut-off, for $k \leq m-1$, we prove convergence with explicit control of error probabilities for each of the $\log \log N$ active components.

We thereby obtain results of the form

$$\begin{aligned} X_t^k &= x_t^k + a_k O(\sqrt{\log \log \log N / Na_k}), \\ X_t^k &= x_t^k + N^{-1/2} \gamma_t^k + a_k O(\log(Na_k) / Na_k). \end{aligned}$$

Note that each component is estimated in the correct scale, with an error depending on the number of queues active at that level. The $\log \log \log N$ in the first equation is a (small) price we pay for working with infinitely many components. These asymptotics will be established with a degree of uniformity in x_0 , which thus allows a dependence of x_0 on N . The Gaussian approximation relies, as in the finite-dimensional case, on a sophisticated coupling of the compensated Poisson process with Brownian motion due to Komlós, Major and Tusnády [8].

We will give a third result, for $k \leq m-1$, of the form

$$X_t^k = x_t^k + N^{-1/2} \tilde{\gamma}_t^k + a_k O((\log \log \log N / Na_k)^{3/4}).$$

Here $(\tilde{\gamma}_t)_{t \geq 0}$ is a jump process with drift which depends on N but is of a simpler type than X in that it is a linear function of additive Poisson noise. The characteristics of $\tilde{\gamma}$ are derived in a simple and canonical way from those of X . Moreover $\tilde{\gamma}$ and X share a common filtration. The error term is larger than for the Gaussian approximation. On the other hand the derivation is significantly simpler.

We obtain also the behaviour of the queue sizes at and above the cut-off. We see a residual randomness in $(X_t^m)_{t \geq 0}$ even for large values of N . This may be approximated in terms of an $M/M/\infty$ queue with arrival rate $N\lambda(x_t^{m-1})^d$ and service rate 1. Over a given finite time interval, there are no queues with lengths greater than m .

Thus we will show for the supermarket model that its infinite-dimensional character does not prevent the derivation of precise asymptotics. We expect the general approach taken here to adapt well to a number of further examples of similar character.

2 Statement of results

Let S_0 denote the set of non-increasing sequences $x = (x^k : k \in \mathbb{N})$ in $[0, 1]$. For $x \in S_0$, set $x^0 = 1$ and define $\lambda_+^k(x) = \lambda((x^{k-1})^d - (x^k)^d)$, $\lambda_-^k(x) = x^k - x^{k+1}$ and $b^k(x) = \lambda_+^k(x) - \lambda_-^k(x)$. It is shown in [15] that, given $x_0 \in S_0$, there is a unique solution $(x_t)_{t \geq 0}$ to $\dot{x}_t = b(x_t)$ in S_0 . Moreover, for any other solution $(y_t)_{t \geq 0}$ in S_0 , $x_0^k \leq y_0^k$ for all k implies $x_t^k \leq y_t^k$ for all k and all $t \geq 0$.

Recall that $a_k = \lambda^{1+d+\dots+d^{k-1}}$ for $k \in \mathbb{N}$. Then $a = (a_k : k \in \mathbb{N})$ is the unique solution in S_0 to $b(x) = 0$ such that $\lim_{k \rightarrow \infty} x^k = 0$; to see this note that $x^{k+1} - \lambda(x^k)^d$ is independent of k for any solution x . Define $\|x\| = \sup_k |x^k|/a_k$, set $E = \{x \in \mathbb{R}^{\mathbb{N}} : \|x\| < \infty\}$ and set $S = S_0 \cap E$. Extend $\lambda_{\pm}$ and b to $\mathbb{R}^{\mathbb{N}}$ by setting $\lambda_+^k(x) = \lambda((y^{k-1})^d - (y^k)^d)^+$, $\lambda_-^k(x) = (y^k - y^{k+1})^+$, where $y^k = (x^k)^+$. It is easy to check that b maps E to itself and is locally Lipschitz for the given norm. Thus, if $x_0 \in S$, then $(x_t)_{t \geq 0}$ does not leave S immediately. Moreover, if $\|x_0\| \leq 1$, then, by comparison with the stationary solution a , $\|x_t\| \leq 1$ for all $t \geq 0$. For $\rho \geq 1$ and $t_0 > 0$ set

$$S(\rho, t_0) = \{x_0 \in S : \|x_t\| \leq \rho \text{ for all } t \leq t_0\}.$$

The state-space I of the Markov chain $X = (X_t)_{t \geq 0}$ is the set of non-increasing sequences in $N^{-1}\{0, 1, \dots, N\}$ with finitely many non-zero terms. Thus $I \subseteq S$. The Lévy kernel for X is given by

$$K(x, dy) = \sum_{k=1}^{\infty} [N\lambda_+^k(x)\delta_{e_k/N}(dy) + N\lambda_-^k(x)\delta_{-e_k/N}(dy)],$$

where e_k denotes the k -th standard basis vector. Given $m \in \mathbb{N}$, let $(\hat{X}_t^m)_{t \geq t_0}$ be a process starting from x_0^m and such that $N\hat{X}^m$ is an $M/M/\infty$ queue, with arrival rate $N\lambda(x_t^{m-1})^d$ and service rate 1. We can now state our law of large numbers.

Theorem 2.1. *Set $m = m(N) = \inf\{k \in \mathbb{N} : Na_k \leq (\log N)^4\}$. There is a coupling of $\hat{X}^m$ and X^m such that, for all $\rho \geq 1, t_0 > 0$ and all sequences $R(N)$ with $R(N)/\sqrt{\log \log \log N} \rightarrow \infty$, we have*

$$\sup_{x_0 \in I \cap S(\rho, t_0)} \mathbb{P}_{x_0}(\sqrt{N}|X_t^{N,k} - x_t^k| > R(N)\sqrt{a_k} \text{ for some } k \leq m-1 \\ \text{or } X_t^{N,m} \neq \hat{X}_t^{N,m} \text{ or } X_t^{N,m+1} \neq 0 \text{ for some } t \leq t_0) \rightarrow 0.$$

In particular

$$\sup_{k \leq m-1} \sup_{t \leq t_0} |X_t^{k,N} - x_t^k|/a_k \rightarrow 0$$

in probability, uniformly in $x_0 \in I \cap S(\rho, t_0)$.

We know, see [15], that if $\rho > 0$ and $\|x_0\| \leq \rho$ then $x_t^k \rightarrow a_k$ as $t \rightarrow \infty$. Thus, for $k \leq m-1$, the proportional error in approximating X_t^k by the deterministic process x_t^k is small for large values of N .

The central limit theorem shows generically that the power $\sqrt{N}$ in Theorem 2.1 cannot be improved while the approximating process $(x_t)_{t \geq 0}$ remains deterministic. Our next result is a refined approximation which allows an improvement to $N^{3/4}$. Let $\tilde{\mu}$ be a Poisson random measure on $\mathbb{R}^{\mathbb{N}} \times (0, t_0]$ with intensity

$$\tilde{\nu}(dy, dt) = K(x_t, dy)dt.$$

We show in Section 6 that the linear equations

$$\tilde{\gamma}_t^k = \sqrt{N} \int_{\mathbb{R}^{\mathbb{N}} \times (0, t]} y^k (\tilde{\mu} - \tilde{\nu})(dy, ds) + \int_0^t \nabla b^k(x_s) \tilde{\gamma}_s ds \quad (1)$$

have a unique cadlag solution $(\tilde{\gamma}_t^k : k \in \mathbb{N}, t \leq t_0)$. Set $\tilde{X}_t = x_t + N^{-1/2} \tilde{\gamma}_t$.

Theorem 2.2. *Define $m(N)$ as in Theorem 2.1. There is a coupling of $\tilde{X}$ and X , in a common filtration, such that, for all $\rho \geq 1, t_0 > 0$ and all sequences $\tilde{R}(N)$ with $\tilde{R}(N)/(\log \log \log N)^{3/4} \rightarrow \infty$, we have*

$$\sup_{x_0 \in I \cap S(\rho, t_0)} \mathbb{P}_{x_0}(N^{3/4} |X_t^k - \tilde{X}_t^k| > \tilde{R}(N) a_k^{1/4} \text{ for some } k \leq m-1, t \leq t_0) \rightarrow 0.$$

The final result is a diffusion approximation. Let $B_+^k, B_-^k, k \in \mathbb{N}$, be independent standard Brownian motions. Set $\sigma_{\pm}^k(x) = \sqrt{\lambda_{\pm}^k(x)}$. We show in Section 6 that the linear equations

$$\gamma_t^k = \int_0^t \sigma_+^k(x_s) dB_+^k(s) - \int_0^t \sigma_-^k(x_s) dB_-^k(s) + \int_0^t \nabla b^k(x_s) \gamma_s ds, \quad t \leq t_0 \quad (2)$$

have a unique solution $(\gamma_t^k : k \in \mathbb{N}, t \leq t_0)$ with

$$\sup_{k \in \mathbb{N}} \mathbb{E}(\sup_{t \leq t_0} |\gamma_t^k|^2) < \infty.$$

Set $\bar{X}_t = x_t + N^{-1/2} \gamma_t$.

Theorem 2.3. *Define $m(N)$ as in Theorem 2.1. There is a coupling of $\bar{X}$ and X such that, for all $\rho \geq 1, t_0 > 0$, there is a constant $\bar{R}$, independent of N , such that*

$$\sup_{x_0 \in I \cap S(\rho, t_0)} \mathbb{P}_{x_0}(N|X_t^k - \bar{X}_t^k| > \bar{R} \log(Na_k) \text{ for some } k \leq m-1, t \leq t_0) \rightarrow 0.$$

We remark that there are alternative versions of all three theorems in which x_0 is replaced by $x_0^{(m)} = (x_0^1, \dots, x_0^{m-1}, 0, \dots)$ and $\lambda_{\pm}(x)$ is replaced by $\lambda_{\pm}(x)^{(m)}$, so that the approximating deterministic dynamics are $(m-1)$ -dimensional. The proofs are a minor modification of the proofs given below. These alternative versions would have merit in any computational implementation of the approximations since m is of order only $\log \log N$.

3 Law of large numbers

In the first half of this section, we fix N and $A, R \geq 1$ and set $m = \inf\{k \in \mathbb{N} : Na_k \leq A\}$. We will obtain, subject to certain constraints, a global estimate on the probability appearing in Theorem 2.1. In the second half we will show that this estimate becomes small as $N \rightarrow \infty$ when $A = (\log N)^4$ and when R is chosen as in Theorem 2.1.

Fix $\rho \geq 1$ and $t_0 > 0$ and assume that $\rho A^d \leq N^{d-1}$. Set $\alpha = (\log \log N - \log \log(1/\lambda))/\log d$, so that $N\lambda^{d\alpha} = 1$ and $Na_k = \lambda^{1+d+\dots+d^{k-1}-d\alpha}$. If $k \leq \alpha$ then $Na_k \geq 1$, whereas if $k \geq \alpha + 1$ then $Na_k \leq 1$. Hence, at least for sufficiently large N , we will have $m \in (\alpha - 1, \alpha + 2)$.

Consider the case $x_0 \in S(\rho, t_0)$. Then $Nx_0^{m+1} \leq N\rho a_{m+1} < N\rho a_m^d \leq \rho A^d/N^{d-1} \leq 1$, so $x_0^{m+1} = 0$. Set $T_1 = \inf\{t \geq 0 : X_t^{m+1} \neq 0\}$. Note that, while $X_t^{m+1} = 0$, X_t^{m+1} increases at rate $N\lambda(X_{t-}^m)^d$, whereas $X_t^m + X_t^{m+1}$ increases at rate $N\lambda(X_{t-}^{m-1})^d$ and decreases at rate NX_{t-}^m . We can therefore find an $M/M/\infty$ queue $(Q_t)_{t \geq 0}$, starting from Nx_0^m , without arrivals and with service rate 1, and a Poisson random measure $\mu(dt, dx, du)$ on $(0, \infty)^3$, independent of Q and of intensity $e^{-u} dt dx du$, such that, for $t \leq T_1$,

$$NX_t^{m+1} = \mu(\{(s, x, u) : s \leq t < s + u, x \leq N\lambda(X_{s-}^m)^d\})$$

and

$$N(X_t^m + X_t^{m+1}) = Q_t + \mu(\{(s, x, u) : s \leq t < s + u, x \leq N\lambda(X_{s-}^{m-1})^d\}).$$

Define $(\hat{X}_t^m)_{t \geq 0}$ by

$$N\hat{X}_t^m = Q_t + \mu(\{(s, x, u) : s \leq t < s + u, x \leq N\lambda(x_s^{m-1})^d\})$$

and set $T_2 = \inf\{t \geq 0 : X_t^m + X_t^{m+1} \neq \hat{X}_t^m\}$. Then $(N\hat{X}_t^m)_{t \geq 0}$ is an $M/M/\infty$ queue starting from Nx_0^m , with arrival rate $N\lambda(x_t^{m-1})^d$ and service rate 1. Fix $r > 1$ and set $T_3 = \inf\{t \geq 0 : \hat{X}_t^m > ra_m\}$. Fix $R \geq 1$, set

$$T_4 = \inf\{t \geq 0 : \sqrt{N}|X_t^k - x_t^k| > R\sqrt{a_k} \text{ for some } k \leq m-1\},$$

and set $T = T_1 \wedge T_2 \wedge T_3 \wedge T_4 \wedge t_0$. Finally, set

$$p = p(N, \lambda, d, x_0, A, R, r) = \mathbb{P}(T < t_0).$$

Proposition 3.1. *Assume that $x_0 \in S(\rho, t_0)$, that $A, R, \rho \geq 1$ with $\rho A^d \leq N^{d-1}$, that $r > \rho$ and that*

$$2rAt_0e^{Lt_0}/N^{(1/2)(1-1/d)} \leq R \leq (t_0 \wedge 1)\sqrt{A},$$

where $L = 2(d\sigma^{d-1} + 1)$ and $\sigma = \rho + 1$. Then $p \leq p_1 + p_2 + p_3 + p_4$, where

$$\begin{aligned} p_1 &= A^d r^d t_0 / N^{d-1}, \\ p_2 &= A^{1-1/(2d)} d\sigma^{d-1} R t_0 / N^{(1/2)(1-1/d)}, \\ p_3 &= \rho^d t_0 / (r - \rho), \\ p_4 &= 2m \exp(-R^2 / (20\sigma^d t_0 e^{2Lt_0})). \end{aligned}$$

Proof. It will suffice to show that $\mathbb{P}(T = T_i) \leq p_i$ for $i = 1, 2, 3, 4$. Recall that for a Poisson random variable Y of parameter $\nu > 0$ and for $a > 0$ we have $\mathbb{P}(Y \geq a) \leq \nu/a$. For $t < T$ we have $N\lambda(X_t^m)^d \leq N\lambda(ra_m)^d \leq A^d r^d / N^{d-1}$, so X_T^{m+1} is dominated by a Poisson random variable Y_1 of parameter p_1 , and so

$$\mathbb{P}(T = T_1) = \mathbb{P}(X_T^{m+1} = 1) \leq \mathbb{P}(Y_1 \geq 1) \leq p_1.$$

Since $x_0 \in S(\rho, t_0)$, we have $x_t^k \leq \rho a_k$ for all $k \in \mathbb{N}$ and all $t \leq t_0$. For $k \leq m-1$ we have $R\sqrt{a_k/N} \leq a_k R / \sqrt{Na_k} \leq a_k R / \sqrt{A} \leq a_k$. Hence, for $t < T$ and $k \leq m-1$,

$$X_t^k \leq x_t^k + R\sqrt{a_k/N} \leq \sigma a_k. \quad (3)$$

Then, for $t < T$,

$$\begin{aligned} N\lambda|(X_t^{m-1})^d - (x_t^{m-1})^d| &\leq N\lambda d\sigma^{d-1}a_{m-1}^{d-1}|X_t^{m-1} - x_t^{m-1}| \\ &\leq N\lambda d\sigma^{d-1}a_{m-1}^{d-1}R\sqrt{a_{m-1}/N} \leq A^{1-1/(2d)}d\sigma^{d-1}R/N^{(1/2)(1-1/d)}. \end{aligned}$$

Set

$$\Delta = \mu(\{(t, x, u) : t \leq T, N\lambda(x_t^{m-1} \wedge X_{t-}^{m-1})^d < x \leq N\lambda(x_t^{m-1} \vee X_{t-}^{m-1})^d\}).$$

Then Δ is dominated by a Poisson random variable Y_2 of parameter p_2 . Hence

$$\mathbb{P}(T = T_2) = \mathbb{P}(\Delta = 1) \leq \mathbb{P}(Y_2 \geq 1) \leq p_2.$$

Note that $\lambda(x_t^{m-1})^d \leq \lambda\rho^d a_{m-1}^d = \rho^d a_m$ for all $t \leq t_0$. Thus $N\hat{X}_T^m \leq Nx_0^m + Y_3$ for a Poisson random variable Y_3 of parameter $N\rho^d a_m t_0$ and so

$$\mathbb{P}(T = T_3) = \mathbb{P}(\hat{X}_T^m \geq ra_m) \leq \mathbb{P}(Y_3 \geq N(r - \rho)a_m) \leq p_3.$$

It remains to estimate $\mathbb{P}(T = T_4)$. For this we write

$$X_t^k = x_0^k + M_t^k + \int_0^t b^k(X_s)ds \quad (4)$$

so that

$$X_t^k - x_t^k = M_t^k + \int_0^t (b^k(X_s) - b^k(x_s))ds.$$

Then we use a combination of exponential martingale inequalities and Gronwall's lemma to obtain the desired estimate. First we investigate how small M will need to be to obtain the required bound on $|X_t^k - x_t^k|$ for $k \leq m-1$. Note that

$$|\lambda_+^k(x) - \lambda_+^k(y)| \leq \lambda d(x^{k-1} \vee y^{k-1})^{d-1}|x^{k-1} - y^{k-1}| + \lambda d(x^k \vee y^k)^{d-1}|x^k - y^k|$$

so, provided that $x^k \vee y^k \leq \sigma a_k$ and $x^{k-1} \vee y^{k-1} \leq \sigma a_{k-1}$,

$$|\lambda_+^k(x) - \lambda_+^k(y)| \leq d\sigma^{d-1}\{(a_k/a_{k-1})|x^{k-1} - y^{k-1}| + |x^k - y^k|\}. \quad (5)$$

Also

$$|\lambda_-^k(x) - \lambda_-^k(y)| \leq |x^k - y^k| + |x^{k+1} - y^{k+1}|. \quad (6)$$

Hence, provided that $x^k \vee y^k \leq \sigma a_k$ and $x^{k-1} \vee y^{k-1} \leq \sigma a_{k-1}$, we have

$$|b^k(x) - b^k(y)|/\sqrt{a_k} \leq L \sup_{j=k-1, k, k+1} |x^j - y^j|/\sqrt{a_j}.$$

We note that the definitions of T_2 and T_3 force $X_t^m \leq ra_m$ for all $t < T$. Set

$$f(t) = \sup_{k \leq m-1} \sup_{s \leq t} |X_s^k - x_s^k|/\sqrt{a_k}.$$

Then, for $t < T$ and $k \leq m-2$,

$$|b^k(X_t) - b^k(x_t)|/\sqrt{a_k} \leq Lf(t)$$

and

$$|b^{m-1}(X_t) - b^{m-1}(x_t)|/\sqrt{a_{m-1}} \leq Lf(t) + ra_m/\sqrt{a_{m-1}}.$$

Hence, for $t \leq T$,

$$f(t) \leq (M_t^* + ra_mt/\sqrt{a_{m-1}}) + L \int_0^t f(s)ds$$

where

$$M_t^* = \sup_{k \leq m-1} \sup_{s \leq t} |M_s^k|/\sqrt{a_k}.$$

Set $\alpha_k = \frac{1}{2}e^{-Lt_0}R\sqrt{a_k/N}$ and consider for $k \leq m-1$ the stopping times $T^k = T_-^k \wedge T_+^k$, where

$$T_{\pm}^k = \inf\{t \geq 0 : \pm M_t > \alpha_k\}.$$

Suppose that $T < T^1 \wedge \dots \wedge T^{m-1}$. Then $M_T^* \leq \frac{1}{2}e^{-Lt_0}R/\sqrt{N}$. On the other hand

$$ra_mt_0/\sqrt{a_{m-1}} = \lambda^{1/(2d)}ra_m^{1-1/(2d)}t_0 \leq rA^{1-1/(2d)}t_0/N^{1-1/(2d)} \leq (1/2)e^{-Lt_0}R/N^{1/2}.$$

So by Gronwall's lemma

$$f(T) \leq e^{Lt_0}(M_T^* + ra_mt_0/\sqrt{a_{m-1}}) \leq R/\sqrt{N}.$$

Hence

$$\mathbb{P}(T = T_4) \leq \sum_{k=1}^{m-1} \mathbb{P}(T^k \leq T)$$

and it remains to estimate $\mathbb{P}(T_{\pm}^k \leq T)$ for $k \leq m-1$. For $k \in \mathbb{N}, x \in S$ and $\theta \in \mathbb{R}$, set

$$\phi^k(x, \theta) = \lambda_+^k(x)h(\theta) + \lambda_-^k(x)h(-\theta),$$

where $h(\theta) = e^\theta - 1 - \theta$. For $t < T$ and $k \leq m-1$ we have $X_t^k \leq \sigma a_k$ so

$$\phi^k(X_t, \theta) \leq \lambda \sigma^d a_{k-1}^d h(\theta) + \sigma a_k h(-\theta) \leq \sigma^d a_k g(\theta) \quad (7)$$

where $g(\theta) = e^\theta - 2 + e^{-\theta}$. Consider for $\theta \geq 0$ the exponential martingale

$$\begin{aligned} Z_t^k &= \exp\{N\theta(X_t^k - X_0^k) - \int_0^t \int_{\mathbb{R}^N} (e^{N\theta y^k} - 1)K(X_s, dy)ds\} \\ &= \exp\{N\theta M_t^k - N \int_0^t \phi^k(X_s, \theta)ds\} \end{aligned}$$

and note that on the event $T_+^k \leq T$ we have

$$Z_{T_+^k}^k \geq \exp\{N\theta\alpha_k - N\sigma^d a_k g(\theta)t_0\}.$$

By optional stopping $\mathbb{E}(Z_{T_+^k}^k) \leq 1$, so by Chebyshev's inequality

$$\mathbb{P}(T_+^k \leq T) \leq \exp(-N\theta\alpha_k + N\sigma^d a_k g(\theta)t_0).$$

We choose $\theta = \alpha_k/(2\sigma^d a_k \gamma t_0)$, where $\gamma = g(1) \leq 5/4$. It is straightforward to check that $\theta \leq 1$ so $g(\theta) \leq \gamma\theta^2$. Hence

$$\mathbb{P}(T_+^k \leq T) \leq \exp(-N\alpha_k^2/(4\sigma^d a_k \gamma t_0)) = \exp(-R^2/(16\sigma^d \gamma t_0 e^{2Lt_0})).$$

The same bound applies to $\mathbb{P}(T_-^k \leq T)$. So we have shown that $\mathbb{P}(T_4 = T) \leq p_4$ as required. $\square$

Proof of Theorem 2.1. We will determine conditions on sequences $R(N)$ and $r(N)$ so that, for $A(N) = (\log N)^4$, as $N \rightarrow \infty$, all the constraints of Proposition 3.1 are satisfied and, with an obvious notation, $p_i(N) \rightarrow 0$ for $i = 1, 2, 3, 4$. For $p_4(N) \rightarrow 0$ it suffices that $\log \log N \exp(-R^2/(20\sigma^d t_0 e^{2Lt_0})) \rightarrow 0$ and hence that $R/\sqrt{\log \log \log N} \rightarrow \infty$. For $p_3(n) \rightarrow 0$ it suffices that $r \rightarrow \infty$. For $p_2(N) \rightarrow 0$ it suffices that $A^{2-1/d} R^2/N^{1-1/d} \rightarrow 0$ and for $p_1(N) \rightarrow 0$ it suffices that $Ar/N^{1-1/d} \rightarrow 0$. If we can also arrange that $R/\sqrt{A} \rightarrow 0$ and $rA/(RN^{(1/2)(1-1/d)}) \rightarrow 0$ then all the constraints of Proposition 3.1 will be satisfied eventually. A possible choice is to take $r(N) =$

$N^{(1/2)(1-1/d)}/(\log N)^4$ and any sequence $R(N)$ with $R(N)/\sqrt{\log \log \log N} \rightarrow \infty$ and $R(N)/(\log N)^2 \rightarrow 0$. This proves the first part of the theorem. For the final assertion it suffices to note that for $k \leq m-1$, $R(N)/\sqrt{Na_k} \leq R(N)/\sqrt{A(N)} \rightarrow 0$. $\square$

We remark that the choice $R(N) = \log N$ leads to a bound of the form $p(N) \leq CN^{-(1/2)(1-1/d)}$ up to logarithmic corrections. This is the best rate of decay of probabilities we have found. We remark also that a marginally shorter proof can be had by replacing the exponential martingale inequality by Doob's L^2 -inequality, at the small cost of requiring that $R(N)/\sqrt{\log \log N} \rightarrow \infty$.

4 A refinement of the fluid limit

This section leads to a proof of Theorem 2.2. The deterministic limit (for components $k \leq m-1$) just discussed will be refined by approximating the martingale M in (4) by another martingale whose characteristics are determined by the limit path, and at the same time linearizing around the limit path. The accuracy of the approximation is thereby improved from $N^{-1/2}$ to $N^{-3/4}$ at the cost of moving to an approximating process which is not deterministic but has a simple random structure, being a linear function of a Poisson random measure.

Define a measure $\tilde{\nu}$ on $\mathbb{R}^N \times (0, t_0]$ by

$$\tilde{\nu}(dy, dt) = K(x_t, dy)dt.$$

We will take $\tilde{\mu}$ to be a Poisson random measure with intensity $\tilde{\nu}$ coupled, in a way to be specified, with the process X . Define $\tilde{M} = (\tilde{M}_t^k : k \in \mathbb{N}, t \leq t_0)$ by

$$\tilde{M}_t^k = \int_{\mathbb{R}^N \times (0, t]} y^k (\tilde{\mu} - \tilde{\nu})(dy, ds)$$

and define $\tilde{\gamma} = (\tilde{\gamma}_t^k : k \in \mathbb{N}, t \leq t_0)$ by

$$\tilde{\gamma}_t = \sqrt{N}\tilde{M}_t + \int_0^t \nabla b(x_s) \tilde{\gamma}_s ds.$$

We show in Section 6 that we can write $\tilde{\gamma}$ as an explicit linear function of $\tilde{\mu} - \tilde{\nu}$

$$\tilde{\gamma}_t = \sqrt{N} \int_{\mathbb{R}^N \times (0, t]} \Phi_{t,s} y (\tilde{\mu} - \tilde{\nu})(dy, ds) \quad (8)$$

where $(\Phi_{t,s} : s \leq t \leq t_0)$ is the $\mathbb{N} \times \mathbb{N}$ matrix-valued process given by

$$\frac{\partial}{\partial t} \Phi_{t,s} = \nabla b(x_t) \Phi_{t,s}, \quad \Phi_{s,s} = I.$$

Thus $\tilde{\gamma}$ has a simpler stochastic structure than X . In particular, we can write the characteristic function of any finite-dimensional distribution of $\tilde{\gamma}$ in terms of $(x_t)_{t \leq t_0}$ and $(\Phi_{t,s} : s \leq t \leq t_0)$.

Recall that

$$X_t = x_0 + M_t + \int_0^t b(X_s) ds.$$

On the other hand, if we set $\tilde{X}_t = x_t + N^{-1/2} \tilde{\gamma}_t$, then

$$\tilde{X}_t = x_0 + \tilde{M}_t + \int_0^t b(x_s) ds + \int_0^t \nabla b(x_s) (\tilde{X}_s - x_s) ds.$$

Set $\tilde{Y} = X - \tilde{X}$, $\tilde{D} = M - \tilde{M}$ and

$$A_t = \int_0^t (b(X_s) - b(x_s) - \nabla b(x_s)(X_s - x_s)) ds. \quad (9)$$

Then

$$\tilde{Y}_t = \tilde{D}_t + A_t + \int_0^t \nabla b(x_s) \tilde{Y}_s ds. \quad (10)$$

We will obtain a good approximation if we can couple $\tilde{M}$ with M to make $\tilde{D}$ small. Define kernels K_0, K_+, K_- on $(0, t_0] \times E \times \mathbb{R}^{\mathbb{N}}$ by

$$\begin{aligned} K_0(t, x, dy) &= K(x, dy) \wedge K(x_t, dy), \\ K_{\pm}(t, x, dy) &= (K(x, dy) - K(x_t, dy))^{\pm}, \end{aligned}$$

and let $K_*(t, x, w, dx', dw')$ be the image of the measure

$$K_0(t, x, dy_0) \otimes K_+(t, x, dy_+) \otimes K_-(t, x, dy_-)$$

by the map $(x', w') = (y_0 + y_+, y_0 + y_-)$. Let $(X_t, W_t)_{t \geq 0}$ be a Markov chain, starting from $(x_0, 0)$, with time-dependent Lévy kernel K_* . Set

$$\tilde{\mu}_t = \sum_{\Delta W_t \neq 0} \delta_{(t, \Delta W_t)}.$$

Then $(X_t)_{t \geq 0}$ is a Markov chain with Lévy kernel K and $\tilde{\mu}$ is a Poisson random measure with intensity $\tilde{\nu}$. We have coupled X and W so that, as far as possible, they have the same jumps. Set $T_5 = \inf\{t \geq 0 : |\tilde{\gamma}_t^m| > r\sqrt{a_m}\}$, where r is as in the previous section. Fix $\tilde{R} > 0$ and set

$$T_6 = \inf\{t \geq 0 : N^{3/4}|X_t^k - \tilde{X}_t^k| > \tilde{R}a_k^{1/4} \text{ for some } k \leq m-1\}.$$

Finally, set $\tilde{T} = T \wedge T_5 \wedge T_6$ and $\tilde{p} = \mathbb{P}(\tilde{T} < t_0)$.

Proposition 4.1. *Assume that the conditions of Proposition 3.1 hold. Set $H = \frac{1}{2}d(d-1)\sigma^{d-2}$ and assume in addition that $rA \leq N^{(1/2)(1-1/d)}$ and*

$$4HR^2t_0e^{Lt_0}/N^{1/4} + 4rAt_0e^{Lt_0}/N^{(1/4)(1-1/d)} \leq \tilde{R} \leq 4RLt_0e^{Lt_0}A^{1/4}.$$

Then $\tilde{p} \leq p_1 + p_2 + p_3 + p_4 + p_5 + p_6$, where p_1, p_2, p_3, p_4 are defined in Proposition 3.1 and

$$p_5 = 8(\rho^d + 1)t_0e^{2Lt_0^2}/r^2, \quad p_6 = 2m \exp\{-\tilde{R}^2/(20RLt_0e^{2Lt_0})\}.$$

Proof. Given Proposition 3.1, it will suffice to show that $\mathbb{P}(\tilde{T} = T_i) \leq p_i$ for $i = 5, 6$. By Proposition 6.1,

$$\mathbb{E}(\sup_{t \leq t_0} |\tilde{\gamma}_t^m|^2) \leq 8(\rho^d + 1)t_0e^{2Lt_0^2}a_m.$$

So $\mathbb{P}(\tilde{T} = T_5) \leq p_5$ by Chebyshev's inequality.

We now follow an argument similar to the proof that $\mathbb{P}(T = T_4) \leq p_4$ in Proposition 3.1. Set

$$\begin{aligned} \tilde{f}(t) &= \sup_{k \leq m-1} \sup_{s \leq t} |\tilde{Y}_s^k|/a_k^{1/4}, \\ \tilde{A}_t^* &= \sup_{k \leq m-1} \sup_{s \leq t} |A_s^k|/a_k^{1/4} + \int_0^t |\tilde{Y}_s^m|ds/a_{m-1}^{1/4}, \\ \tilde{D}_t^* &= \sup_{k \leq m-1} \sup_{s \leq t} |\tilde{D}_s^k|/a_k^{1/4}. \end{aligned}$$

We note that

$$\nabla b^k(x)y = \lambda d(x^{k-1})^{d-1}y^{k-1} - \lambda d(x^k)^{d-1}y^k - y^k + y^{k+1}$$

so, provided that $x^k \leq \sigma a_k$ and $x^{k-1} \leq \sigma a_{k-1}$,

$$|\nabla b^k(x)y|/a_k^{1/4} \leq L \sup_{j=k-1, k, k+1} |y^j|/a_j^{1/4}.$$

For $t \leq t_0$ we have $x_t^k \leq \rho a_k \leq \sigma a_k$ for all k , so for $k \leq m-1$

$$|\nabla b^k(x_t) \tilde{Y}_t|/a_k^{1/4} \leq L \tilde{f}(t) + \delta_{k,m-1} |\tilde{Y}_t^m|/a_{m-1}^{1/4}.$$

Then, from equation (10) we get

$$\tilde{f}(t) \leq \tilde{D}_t^* + \tilde{A}_t^* + L \int_0^t \tilde{f}(s) ds,$$

so, by Gronwall's lemma, $\tilde{f}(t) \leq e^{Lt}(\tilde{A}_t^* + \tilde{D}_t^*)$ for all $t \leq t_0$. Note that, for $k \leq m-1$,

$$\begin{aligned} b^k(y) - b^k(x) - \nabla b^k(x)(y-x) &= \lambda((y^{k-1})^d - (x^{k-1})^d - d(x^{k-1})^{d-1}(y^{k-1} - x^{k-1})) \\ &\quad - \lambda((y^k)^d - (x^k)^d - d(x^k)^{d-1}(y^k - x^k)) \end{aligned}$$

so, provided that $x^k, y^k \leq \sigma a_k$ and $x^{k-1}, y^{k-1} \leq \sigma a_{k-1}$,

$$\begin{aligned} |b^k(y) - b^k(x) - \nabla b^k(x)(y-x)| &\leq H\lambda(a_{k-1}^{d-2}|y^{k-1} - x^{k-1}|^2 + a_k^{d-2}|y^k - x^k|^2). \end{aligned}$$

For $t < \tilde{T}$ and $k \leq m-1$ we have $|X_t^k - x_t^k| \leq R\sqrt{a_k/N}$; moreover, as we showed at (3), this implies that $X_t^k \leq \sigma a_k$. Hence

$$\begin{aligned} |b^k(X_t) - b^k(x_t) - \nabla b^k(x_t)(X_t - x_t)|/a_k^{1/4} &\leq H\lambda R^2(a_{k-1}^{d-1} + a_k^{d-1})/(Na_k^{1/4}) \leq 2CR^2/N. \end{aligned}$$

Also, $|\tilde{Y}_t^m| \leq |X_t^m - x_t^m| + N^{-1/2}|\tilde{\gamma}_t^m|$ so, for $t < \tilde{T}$, $|\tilde{Y}_t^m|/a_{m-1}^{1/4} \leq (ra_m + N^{-1/2}r\sqrt{a_m})/a_{m-1}^{1/4} \leq 2r(A/N)^{1-1/(4d)}$. It follows that

$$\tilde{A}_{\tilde{T}}^* \leq 2HR^2t_0/N + 2rt_0(A/N)^{1-1/(4d)} \leq \frac{1}{2}e^{-Lt_0}\tilde{R}/N^{3/4}.$$

Set $\tilde{\alpha}_k = \frac{1}{2}e^{-Lt_0}\tilde{R}a_k^{1/4}/N^{3/4}$ and consider the stopping times $\tilde{T}^k = \tilde{T}_+^k \wedge \tilde{T}_-^k$, where

$$\tilde{T}_\pm^k = \inf\{t \geq 0 : \pm \tilde{D}_t^k > \tilde{\alpha}_k\}.$$

Suppose that $\tilde{T} < \tilde{T}^1 \wedge \dots \wedge \tilde{T}^{m-1}$. Then $\tilde{D}_{\tilde{T}}^* \leq \frac{1}{2}e^{-Lt_0}\tilde{R}/N^{3/4}$ so $f(\tilde{T}) \leq \tilde{R}/N^{3/4}$ and $\tilde{T} < T_6$. Hence

$$\mathbb{P}(\tilde{T} = T_6) \leq \sum_{k=1}^{m-1} \mathbb{P}(\tilde{T}^k \leq \tilde{T})$$

and it remains to estimate $\mathbb{P}(\tilde{T}_\pm^k \leq \tilde{T})$ for $k \leq m-1$.

For $k \leq m-1$ set

$$\begin{aligned}\psi^k(t, x, \theta) = & (\lambda_+^k(x) - \lambda_+^k(x_t))^+ h(\theta) + (\lambda_-^k(x) - \lambda_-^k(x_t))^+ h(-\theta) \\ & + (\lambda_+^k(x) - \lambda_+^k(x_t))^- h(-\theta) + (\lambda_-^k(x) - \lambda_-^k(x_t))^- h(\theta).\end{aligned}$$

Fix $\theta \geq 0$ and consider for $k \leq m-1$ the exponential martingale

$$\begin{aligned}\tilde{Z}_t^k = & \exp\{N\theta(X_t^k - X_0^k - W_t^k) - \int_0^t \int_{\mathbb{R}^N} (e^{N\theta y^k} - 1)K^+(s, X_s, dy)ds \\ & - \int_0^t \int_{\mathbb{R}^N} (e^{-N\theta y^k} - 1)K^-(s, X_s, dy)ds\} \\ = & \exp\{N\theta \tilde{D}_t^k - N \int_0^t \psi^k(s, X_s, \theta)ds\}.\end{aligned}$$

For $k \leq m-2$ and $t \leq t_0$, for $x^k \leq \sigma a_k$ and $x^{k-1} \leq \sigma a_{k-1}$, we can estimate as at (5), (6), (7) to obtain

$$\begin{aligned}\psi^k(t, x, \theta)/\sqrt{a_k} \leq & g(\theta)(|\lambda_+^k(x) - \lambda_+^k(x_t)| + |\lambda_-^k(x) - \lambda_-^k(x_t)|)/\sqrt{a_k} \\ \leq & Lg(\theta) \sup_{j=k-1, k, k+1} |x^j - x_t^j|/\sqrt{a_j}\end{aligned}$$

so, for $t < \tilde{T}$ and $k \leq m-2$,

$$\psi^k(t, X_t, \theta) \leq Lg(\theta)R\sqrt{a_k/N}. \quad (11)$$

Similarly, since we assume $R \geq 1, rA \leq N^{(1/2)(1-1/d)}$ and $Na_m \leq A$, we have, for $t < \tilde{T}$, $X_t^m \leq ra_m \leq R\sqrt{a_{m-1}/N}$ and we can show that (11) remains true for $k = m-1$. By optional stopping we have $\mathbb{E}(\tilde{Z}_{\tilde{T}_+^k}^k) \leq 1$ for all $k \leq m-1$.

But on the event $\tilde{T}_+^k \leq \tilde{T}$ we have

$$\tilde{Z}_{\tilde{T}_+^k}^k \geq \exp\{N\theta \tilde{\alpha}^k - Lg(\theta)Rt_0\sqrt{Na_k}\}.$$

We choose $\theta = 2\sqrt{N}\tilde{\alpha}^k/(5LRt_0\sqrt{a_k})$, checking that $\theta \leq 1$, so that $g(\theta) \leq 5\theta^2/4$, and use Chebyshev's inequality to deduce

$$\mathbb{P}(\tilde{T}_+^k \leq \tilde{T}) \leq \exp\{-N^{3/2}\tilde{\alpha}_k^2/(5LRt_0\sqrt{a_k})\} = \exp\{-\tilde{R}^2/(20LRt_0e^{2Lt_0})\}.$$

The same bound applies to $\mathbb{P}(\tilde{T}_-^k \leq \tilde{T})$. So we have shown that $\mathbb{P}(\tilde{T} = T_6) \leq p_6$ as required. $\square$

Proof of Theorem 2.2. Choose $r(N)$ as in the proof of Theorem 2.1, so that $r(N) \rightarrow \infty$ and so $p_5(N) \rightarrow 0$. Assume $\tilde{R}(N)/(\log N)^2 \rightarrow 0$, and set $s(N) = \tilde{R}(N)/(\log \log \log N)^{3/4}$. Then $s(N) \rightarrow \infty$. Set $R(N) = s(N)(\log \log \log N)^{1/2}$ and $r(N) = N^{(1/4)(1-1/d)}/(\log N)^4$. It is straightforward to check that all the constraints in Propositions 3.1 and 4.1 are satisfied eventually. Moreover, as in the proof of Theorem 2.1, we have $p_i(N) \rightarrow 0$ for $i = 1, 2, 3, 4$. Finally $\tilde{R}(N)^2/R \log \log \log N \rightarrow \infty$ so also $p_6(N) \rightarrow 0$ which proves the theorem. $\square$

5 Diffusion approximation

In this section we prove Theorem 2.3. The method follows the lines set out in [4], Chapter 11. As we have already seen, our process X has around $\log \log N$ active components, which have a wide range of scales. This will require special consideration in the implementation of the general method. We also have to deal with the fact that the variance of the diffusion approximation has degeneracies. The diffusion coefficient, obtained as the square root of the variance, then fails to be Lipschitz and some special care is needed to arrive at the desired convergence.

Let $(X_t^k : k \in \mathbb{N}, t \geq 0)$ be the supermarket process starting from x_0 and recall equation (4)

$$X_t^k = x_0^k + M_t^k + \int_0^t b^k(X_s) ds.$$

Recall also that we set $\bar{X}_t = x_t + N^{-1/2}\gamma_t$, where $(\gamma_t^k : k \in \mathbb{N}, t \leq t_0)$ is defined by the linear equations (2)

$$\gamma_t^k = \sqrt{N}\bar{M}_t^k + \int_0^t \nabla b^k(x_s)\gamma_s ds.$$

and

$$\sqrt{N}\bar{M}_t^k = \int_0^t \sigma_+^k(x_s) dB_+^k(s) - \int_0^t \sigma_-^k(x_s) dB_-^k(s).$$

Set $Y = X - \bar{X}$ and $D = M - \bar{M}$. Then

$$Y_t = D_t + A_t + \int_0^t \nabla b^k(x_s) Y_s ds$$

where A_t is defined at (9). We will obtain a good approximation if we can couple $\bar{M}$ with M to make D small.

The coupling relies on the following approximation result of [8]: there exists a constant $c \in (0, \infty)$ and a probability space on which are defined a compensated Poisson process Z of rate 1 and a standard Brownian motion W such that, for all $t \geq 0$ and $x \in \mathbb{R}$,

$$\mathbb{P}(\sup_{s \leq t} |Z(s) - W(s)| \geq c \log t + x) \leq ce^{-x/c}. \quad (12)$$

Given independent compensated Poisson processes $Z_+^k, Z_-^k, k \in \mathbb{N}$ of rate 1 we can construct X by the equations (4) and

$$M_t^k = N^{-1} \{ Z_+^k (N \int_0^t \lambda_+^k(X_s) ds) - Z_-^k (N \int_0^t \lambda_-^k(X_s) ds) \}.$$

On the other hand, by a theorem of Knight, see for example [14], there exist independent Brownian motions $W_+^k, W_-^k, k \in \mathbb{N}$, such that, for all $k \in \mathbb{N}$ and $t \leq t_0$,

$$W_\pm^k (N \int_0^t \lambda_\pm^k(\bar{X}_s) ds) = \sqrt{N} \int_0^t \sigma_\pm^k(\bar{X}_s) dB_\pm^k(s).$$

The law of $(B_+^k, B_-^k : k \in \mathbb{N})$ given $(W_+^k, W_-^k : k \in \mathbb{N})$ is given by a measurable kernel. So we may assume that these processes are defined on the same probability space as $(Z_+^k, Z_-^k : k \in \mathbb{N})$ and that $(Z_+^k, W_+^k), (Z_-^k, W_-^k), k \in \mathbb{N}$, are independent copies of (Z, W) .

Set $T_7 = \inf\{t \geq 0 : |\gamma_t^m| > r\sqrt{a_m}\}$. Fix $\bar{R} > 0$ and set

$$T_8 = \inf\{t \geq 0 : N|X_t^k - \bar{X}_t^k| > \bar{R} \log(Na_k) \text{ for some } k \leq m-1\}.$$

Finally, set $\bar{T} = T \wedge T_7 \wedge T_8$ and $\bar{p} = \mathbb{P}(\bar{T} < t_0)$.

Proposition 5.1. *Assume that the conditions of Proposition 3.1 hold, and assume moreover that $A \geq e^2$ and $R \leq \sqrt{A}/2$. There is a constant $C < \infty$, depending only on d, λ, ρ and t_0 such that, if*

$$C + C(R^2 + rA)/\log N \leq \bar{R} \leq A/(2 \log A)$$

then $\bar{p} \leq p_1 + p_2 + p_3 + p_4 + p_7 + p_8$, where p_1, p_2, p_3, p_4 are defined in Proposition 3.1 and

$$p_7 = C/r^2, \quad p_8 = Cm(A^{-1} + (\bar{R} \log A)^{-2}).$$

Proof. Given Proposition 3.1, it will suffice to show that $\mathbb{P}(\bar{T} = T_i) \leq p_i$ for $i = 7, 8$. By Proposition 6.1,

$$\mathbb{E}(\sup_{t \leq t_0} |\gamma_t^m|^2) \leq 8(\rho^d + 1)t_0 e^{2Lt_0^2} a_m.$$

So $\mathbb{P}(\bar{T} = T_7) \leq p_7$ for a suitably large C by Chebyshev's inequality. Set

$$\begin{aligned} \bar{f}(t) &= \sup_{k \leq m-1} \sup_{s \leq t} |Y_s^k| / \log(Na_k), \\ A_t^* &= \sup_{k \leq m-1} \sup_{s \leq t} |A_s^k| / \log(Na_k) + \int_0^t |Y_s^m| ds / \log(Na_{m-1}), \\ D_t^* &= \sup_{k \leq m-1} \sup_{s \leq t} |D_s^k| / \log(Na_k). \end{aligned}$$

Since $Na_{m-1} > A \geq e$, we have $Na_k / \log(Na_k) \leq Na_{k-1} / \log(Na_{k-1})$ for all $k \leq m-1$. So we can use an argument from the proof of Proposition 4.1 to obtain $\bar{f}(t) \leq e^{Lt}(A_t^* + D_t^*)$ for all $t \leq t_0$.

The function $f(s) = \log s / (N^{1-d} \lambda s^d)$ is decreasing when $s > e$ and $e < A < Na_{m-1} \leq (AN^{d-1}/\lambda)^{1/d}$, so

$$\log(Na_{m-1}) / (Na_m) = f(Na_{m-1}) \geq f((AN^{d-1}/\lambda)^{1/d}) \geq \log N / (2A).$$

Similarly, $\log(Na_{m-1}) / \sqrt{Na_m} \geq \log N / (2\sqrt{A})$. Hence

$$r(Na_m + \sqrt{Na_m}) / \log(Na_{m-1}) \leq 4rA / \log N.$$

We estimate as in the proof of Proposition 4.1 to obtain for $t < \bar{T}$ and $k \leq m-1$

$$\begin{aligned} &|b^k(X_t) - b^k(x_t) - \nabla b^k(x_t)(X_t - x_t)| / \log(Na_k) \\ &\leq H\lambda R^2(a_{k-1}^{d-1} + a_k^{d-1}) / N \log(Na_k) \leq 2HR^2 / (N \log N) \end{aligned}$$

and

$$|Y_t^m| / \log(Na_{m-1}) \leq (ra_m + N^{-1/2}r\sqrt{a_m}) / \log(Na_{m-1}) \leq 4rA / (N \log N).$$

So $A_{\bar{T}}^* \leq 2t_0(HR^2 + 2rA) / (N \log N) \leq \frac{1}{2}e^{-Lt_0}\bar{R}/N$, provided C is chosen suitably large. For $k \leq m-1$, set $\alpha_k = \frac{1}{2}e^{-Lt_0}\bar{R} \log(Na_k) / N$ and consider

the event $\Omega_k = \{\sup_{t \leq \bar{T}} |D_t^k| > \alpha_k\}$. On $\Omega_0 = \Omega \setminus (\Omega_1 \cup \dots \cup \Omega_{m-1})$ we have $D_{\bar{T}}^* \leq \frac{1}{2}e^{-L t_0} \bar{R}/N$, so $\bar{f}(\bar{T}) \leq \bar{R}/N$ and $\bar{T} < T_8$. Hence

$$\mathbb{P}(\bar{T} = T_8) \leq \sum_{k=1}^{m-1} \mathbb{P}(\Omega_k)$$

and it will suffice to estimate $\mathbb{P}(\Omega_k)$ for each $k \leq m-1$. Fix $k \leq m-1$. We can write $D_t^k = D_+(t) - D_-(t)$ where $D_{\pm}(t) = D_{\pm}^1(t) + D_{\pm}^2(t) + D_{\pm}^3(t)$ and

$$\begin{aligned} D_{\pm}^1(t) &= N^{-1}(Z_{\pm}^k - W_{\pm}^k)(N \int_0^t \lambda_{\pm}^k(X_s) ds) \\ D_{\pm}^2(t) &= N^{-1}\{W_{\pm}^k(N \int_0^t \lambda_{\pm}^k(X_s) ds) - W_{\pm}^k(N \int_0^t \lambda_{\pm}^k(\bar{X}_s) ds)\} \\ D_{\pm}^3(t) &= N^{-1/2} \int_0^t \{\sigma_{\pm}^k(\bar{X}_s) - \sigma_{\pm}^k(x_s)\} dB_{\pm}^k(s). \end{aligned}$$

Hence $\mathbb{P}(\Omega_k) \leq q_+^1 + q_-^1 + q_+^2 + q_-^2 + q_+^3 + q_-^3$ where, for $j = 1, 2, 3$, $q_{\pm}^j = \mathbb{P}(\Omega_{\pm}^j)$ and $\Omega_{\pm}^j = \{\sup_{t \leq \bar{T}} |D_{\pm}^j(t)| > \alpha_k/6\}$.

For $t < \bar{T}$, we have $\lambda_{\pm}^k(X_t) \leq \sigma^d a_k$ so, taking $t = N\sigma^d a_k t_0$ and $x = N\alpha_k/6 - c \log t$ in (12), we obtain

$$\begin{aligned} q_{\pm}^1 &\leq \mathbb{P}\left(\sup_{t \leq N\sigma^d a_k t_0} |Z(t) - W(t)| > N\alpha_k/6\right) \\ &\leq c\sigma^d t_0 N a_k e^{-N\alpha_k/(6c)} = c\sigma^d t_0 N a_k e^{-\bar{R} \log(Na_k)/(12ce^{L t_0})} \\ &= c\sigma^d t_0 (Na_k)^{1-\bar{R}/(12ce^{L t_0})} \leq c\sigma^d t_0 A^{1-\bar{R}/(12ce^{L t_0})} \leq C/(4A), \end{aligned}$$

for a suitable choice of C .

We turn to estimate $q_{\pm}^2$. This will rely on the following continuity estimate for Brownian motion: for $\tau, h, \delta > 0$, setting $n = \lfloor \tau/h \rfloor$,

$$\begin{aligned} &\mathbb{P}\left(\sup_{s, t \leq \tau, |s-t| \leq h} |W(t) - W(s)| > \delta\right) \\ &\leq \mathbb{P}\left(\sup_{k \in \{0, 1, \dots, n-1\}, t \leq 2h} |W(kh+t) - W(kh)| > \delta/2\right) \\ &\leq 2n\mathbb{P}(\sup_{t \leq 2h} W(t) > \delta/2) \leq (2\tau/h)e^{-\delta^2/(16h)}. \end{aligned}$$

For $t < \bar{T}$ we have

$$\begin{aligned} \bar{X}_t^k &\leq x_t^k + |X_t^k - x_t^k| + |\bar{X}_t^k - X_t^k| \leq \rho a_k + R\sqrt{a_k/N} + \bar{R} \log(Na_k)/N \\ &\leq (\rho + R/\sqrt{A} + \bar{R} \log A/A) a_k \leq \sigma a_k, \end{aligned}$$

so

$$N \int_0^t \lambda_{\pm}^k(\bar{X}_s) ds \leq \sigma^d t_0 N a_k.$$

Also, using the inequalities (5) and (6), for $t < \bar{T}$,

$$\begin{aligned} |N \int_0^t (\lambda_+^k(X_s) - \lambda_+^k(\bar{X}_s)) ds| \\ \leq N d \sigma^{d-1} t_0 \bar{R} a_k \{ \log(N a_{k-1}) / (N a_{k-1}) + \log(N a_k) / (N a_k) \} \\ \leq 2 d \sigma^{d-1} t_0 \bar{R} \log(N a_k) \end{aligned}$$

and

$$\begin{aligned} |N \int_0^t (\lambda_-^k(X_s) - \lambda_-^k(\bar{X}_s)) ds| \\ \leq \bar{R} t_0 \{ \log(N a_k) + \log(N a_{k+1}) 1_{k \leq m-2} \} + r t_0 (N a_m + \sqrt{N a_m}) \delta_{k, m-1} \\ \leq 2 t_0 \bar{R} \log(N a_k), \end{aligned}$$

provided that C is sufficiently large. We take $\tau = \sigma^d t_0 N a_k$, $h = 2 d \sigma^{d-1} t_0 \bar{R} \log(N a_k)$ and

$$\delta = N \alpha_k / 6 = \bar{R} \log(N a_k) / (12 e^{L t_0})$$

to obtain

$$q_{\pm}^2 \leq \frac{\sigma N a_k}{d \bar{R} \log(N a_k)} \exp \left\{ - \frac{\bar{R} \log(N a_k)}{4608 d \sigma^{d-1} t_0 e^{2 L t_0}} \right\} \leq C / (4A)$$

for a suitable choice of C .

It remains to estimate $q_{\pm}^3$. We shall show below that there exists a constant C_0 such that, for $t \leq t_0$ and all $k \in \mathbb{N}$,

$$\mathbb{E}(|\gamma_t^k|^2) \leq C_0 (x_t^{k-1} - x_t^k) \wedge (x_t^k - x_t^{k+1}).$$

Then, for $t < \bar{T}$,

$$\begin{aligned} \sqrt{N} |\sigma_-^k(\bar{X}_t) - \sigma_-^k(x_t)| &= \sqrt{N} \left| \sqrt{((\bar{X}_t^k)^+ - (\bar{X}_t^{k+1})^+)^+} - \sqrt{x_t^k - x_t^{k+1}} \right| \\ &= \frac{\sqrt{N} |((\bar{X}_t^k)^+ - (\bar{X}_t^{k+1})^+)^+ - (x_t^k - x_t^{k+1})|}{\sqrt{((\bar{X}_t^k)^+ - (\bar{X}_t^{k+1})^+)^+} + \sqrt{x_t^k - x_t^{k+1}}} \leq (|\gamma_t^k| + |\gamma_t^{k+1}|) / \sqrt{x_t^k - x_t^{k+1}} \end{aligned}$$

so by Doob's L^2 -inequality

$$\mathbb{E}(\sup_{t \leq \bar{T}} |D_-^3(t)|^2) \leq 4\mathbb{E} \int_0^{t_0} N^{-1} |\sigma_-^k(\bar{X}_s) - \sigma_-^k(x_s)|^2 ds \leq 16t_0 C_0 / N^2.$$

Hence

$$q_-^3 = \mathbb{P}(\sup_{t \leq \bar{T}} |D_-^3(t)| > \alpha_k/6) \leq 2304t_0 e^{2Lt_0} C_0 / (\bar{R} \log(Na_k))^2 \leq C(\bar{R} \log A)^{-2}/2$$

for a suitable choice of C .

The argument for q_+^3 is similar. For $t < \bar{T}$,

$$\begin{aligned} & \sqrt{N} |\sigma_+^k(\bar{X}_t) - \sigma_+^k(x_t)| \\ &= \sqrt{N} \left| \sqrt{(\lambda((\bar{X}_t^{k-1})^+)^d - \lambda((\bar{X}_t^k)^+)^d)^+} - \sqrt{\lambda(x_t^{k-1})^d - \lambda(x_t^k)^d} \right| \\ &\leq \lambda d \{ (x_t^{k-1} \vee \bar{X}_t^{k-1})^{d-1} |\gamma_t^{k-1}| + (x_t^k \vee \bar{X}_t^k)^{d-1} |\gamma_t^k| \} / \sqrt{\lambda(x_t^{k-1})^d - \lambda(x_t^k)^d} \\ &\leq \lambda d 2^{d-1} ((x_t^{k-1})^{d-1} + (N^{-1/2} |\gamma_t^{k-1}|)^{d-1}) |\gamma_t^{k-1}| / \sqrt{\lambda(x_t^{k-1} - x_t^k)(x_t^{k-1})^{d-1}} \\ &\quad + \lambda d 2^{d-1} ((x_t^k)^{d-1} + (N^{-1/2} |\gamma_t^k|)^{d-1}) |\gamma_t^k| / \sqrt{\lambda(x_t^{k-1} - x_t^k)(x_t^k)^{d-1}}. \end{aligned}$$

so, by Doob's L^2 -inequality,

$$\begin{aligned} \mathbb{E}(\sup_{t \leq \bar{T}} |D_+^3(t)|^2) &\leq 4\mathbb{E} \int_0^{t_0} N^{-1} |\sigma_+^k(\bar{X}_s) - \sigma_+^k(x_s)|^2 ds \\ &\leq 8d^2 2^{2(d-1)} t_0 (1 + N^{(d-1)/2} C(d)) C_0 / N^2 \end{aligned}$$

where $C(d) = \mathbb{E}(W(1)^{2d})$. Hence

$$\begin{aligned} q_+^3 &= \mathbb{P}(\sup_{t \leq \bar{T}} |D_+^3(t)| > \alpha_k/6) \\ &\leq 2152d^2 2^{2(d-1)} t_0 (1 + N^{(d-1)/2} C(d)) C_0 e^{2Lt_0} / (\bar{R} \log(Na_k))^2 \\ &\leq C(\bar{R} \log A)^{-2}/2 \end{aligned}$$

for a suitable choice of C . On combining this with the bounds for $q_\pm^1$ and $q_\pm^2$ already found, we obtain the desired bound for p_8 . $\square$

Proof of Theorem 2.3. Set $A(N) = r(N) = (\log N)^{1/2}$ and define $\bar{m}(N) = \inf\{k \in \mathbb{N} : Na_k \leq A(N)\}$. Set $R(N) = (\log N)^{1/4}(1 \wedge t_0)/2$. It is straightforward to check that, if C is the constant appearing in Proposition 5.1 and if $\bar{R} = 3C$, then all the constraints in Propositions 3.1 and 5.1 are satisfied eventually and moreover that $p_i(N) \rightarrow 0$ for $i = 1, 2, 3, 4, 7, 8$. Since $\bar{m}(N) \geq m(N)$, this proves the theorem. $\square$

6 Fluctuation variance estimates

We have deferred from other sections the analysis of certain linear equations associated with our processes. The basic questions of existence and uniqueness in suitable spaces may be resolved by standard methods, so we review this only briefly. The more delicate result, Proposition 6.1, which is needed for the diffusion approximation, relies on the particular structure of our model.

We recall the $\mathbb{N} \times \mathbb{N}$ matrix-valued equation

$$\frac{\partial}{\partial t} \Phi_{t,s} = \nabla b(x_t) \Phi_{t,s}, \quad \Phi_{s,s} = I$$

to be solved for $0 \leq s \leq t \leq t_0$. Note that, for $x_0 \in S(\rho, t_0)$ and $t \in [0, t_0]$, we have $\|\nabla b(x_t)\| \leq L$, where $\|\dots\|$ is the operator norm corresponding to $\|x\| = \sup_k |x^k|/a_k$. Hence it is standard that this equation has a unique continuous solution with $\|\Phi_{t,s}\| \leq e^{L(t-s)}$ for all s, t .

The other relevant equations may be considered as stochastic perturbations of the preceding equation. In Theorem 2.2 we used the equation (1)

$$\tilde{\gamma}_t^k = \sqrt{N} \tilde{M}_t^k + \int_0^t \nabla b^k(x_s) \tilde{\gamma}_s ds, \quad t \leq t_0$$

and in Theorem 2.3 we used the equation (2)

$$\gamma_t^k = \sqrt{N} \bar{M}_t^k + \int_0^t \nabla b^k(x_s) \gamma_s ds, \quad t \leq t_0. \quad (13)$$

Here

$$\tilde{M}_t^k = \int_{\mathbb{R}^{\mathbb{N}} \times (0, t]} y^k (\tilde{\mu} - \tilde{\nu})(dy, ds)$$

and

$$\sqrt{N} \bar{M}_t^k = \int_0^t \sigma_+^k(x_s) dB_+^k(s) - \int_0^t \sigma_-^k(x_s) dB_-^k(s).$$

Note that

$$\begin{aligned} N\mathbb{E}(|\tilde{M}_t^k|^2) &= N \int_0^t \int_{\mathbb{R}^N} (y^k)^2 K(x_s, dy) ds \\ &= \int_0^t (\lambda_+^k(x_s) + \lambda_-^k(x_s)) ds = N\mathbb{E}(|M_t^k|^2) \end{aligned}$$

and $\lambda_+^k(x_t) + \lambda_-^k(x_t) \leq (\rho^d + 1)a_k$ for $t \leq t_0$. Hence a standard type of iteration argument, using Doob's L^2 -inequality, shows that these equations have unique measurable solutions with, respectively,

$$\mathbb{E}(\sup_{t \leq t_0} |\tilde{\gamma}_t^k|^2) \leq 8(\rho^d + 1)t_0 e^{2Lt_0^2} a_k$$

and

$$\mathbb{E}(\sup_{t \leq t_0} |\gamma_t^k|^2) \leq 8(\rho^d + 1)t_0 e^{2Lt_0^2} a_k.$$

The details for equation (1) follow below; equation (2) may be treated in the same way. Note that for any vector y and for all $k \in \mathbb{N}$

$$\frac{|\nabla b^k(x_s)y^k|}{\sqrt{a_k}} \leq L \sum_{j=k-1, k, k+1} \frac{|y_j|}{\sqrt{a_j}}.$$

Now let

$$\tilde{\gamma}_t^{(0)} = \sqrt{N}\tilde{M}_t,$$

and for $n \in \mathbb{N}$

$$\tilde{\gamma}_t^{(n)} = \sqrt{N}\tilde{M}_t + \int_0^t \nabla b^k(x_s) \tilde{\gamma}_s^{(n-1)} ds.$$

Then for each k

$$\begin{aligned} \frac{|\tilde{\gamma}_t^{(n+1),k} - \tilde{\gamma}_t^{(n),k}|^2}{a_k} &\leq L^2 \left(\int_0^t \sum_{j=k-1, k, k+1} \frac{|\tilde{\gamma}_s^{(n),j} - \tilde{\gamma}_s^{(n-1),j}|}{\sqrt{a_j}} ds \right)^2 \\ &\leq 3L^2 \sum_{j=k-1, k, k+1} \left(\int_0^t \frac{|\tilde{\gamma}_s^{(n),j} - \tilde{\gamma}_s^{(n-1),j}|}{\sqrt{a_j}} ds \right)^2, \end{aligned}$$

so using Cauchy-Schwartz

$$\frac{|\tilde{\gamma}_t^{(n+1),k} - \tilde{\gamma}_t^{(n),k}|^2}{a_k} \leq 3L^2 t \sum_{j=k-1, k, k+1} \int_0^t \frac{|\tilde{\gamma}_s^{(n),j} - \tilde{\gamma}_s^{(n-1),j}|^2}{a_j} ds.$$

Then for all $0 \leq s \leq t$

$$\sup_{s \leq t} \frac{|\tilde{\gamma}_s^{(n+1),k} - \tilde{\gamma}_s^{(n),k}|^2}{a_k} \leq 3L^2 t \sum_{j=k-1,k,k+1} \int_0^t \sup_{u \leq s} \frac{|\tilde{\gamma}_u^{(n),j} - \tilde{\gamma}_u^{(n-1),j}|^2}{a_j} ds.$$

Let $h^{(n)}(t) = \sup_k \mathbb{E}(\sup_{s \leq t} |\tilde{\gamma}_s^{(n+1),k} - \tilde{\gamma}_s^{(n),k}|^2 / a_k)$; then for $t \leq t_0$

$$h^{(n)}(t) \leq 9L^2 t_0 \int_0^t h^{(n-1)}(s) ds.$$

Hence

$$h^{(n)}(t) \leq \frac{2^n n! (3Lt_0)^{2n}}{(2n)!} \tilde{M}(t_0),$$

where

$$\begin{aligned} \tilde{M}(t) &= \sup_k \mathbb{E}(\sup_{s \leq t} |\tilde{M}_s^k|^2 / a_k) \\ &\leq 4 \sup_k \sup_{s \leq t} \mathbb{E}(|\tilde{M}_s^k|^2 / a_k) \leq 4t(\rho^d + 1). \end{aligned}$$

We deduce that $\tilde{\gamma}_t^{(n)}$ converges to a process $\tilde{\gamma}_t$ uniformly on $[0, t_0]$, and that $\tilde{\gamma}_t$ satisfies (1). The uniqueness part of the proof is similar. Let

$$g(t) = \sup_k \mathbb{E}(\sup_{s \leq t} |\tilde{\gamma}_s^k|^2) / a_k.$$

Then it follows from the above estimates that for $t \leq t_0$

$$g(t) \leq \tilde{M}(t_0) e^{9L^2 t_0^2}.$$

It may be verified by substitution that the formula (8) gives an explicit representation of the solution of equation (1).

Proposition 6.1. *The solution $(\gamma_t^k : k \in \mathbb{N}, t \leq t_0)$ to (2) satisfies*

$$\sup_{k \in \mathbb{N}} \sup_{t \leq t_0} \mathbb{E}(|\gamma_t^k|^2) / \min\{x_t^{k-1} - x_t^k, x_t^k - x_t^{k+1}\} < \infty.$$

Proof. Note that $\lambda_+^k(x_t) \leq d(x_t^{k-1} - x_t^k)$, so

$$\begin{aligned} \dot{x}_t^k - \dot{x}_t^{k+1} &= \lambda_+^k(x_t) - \lambda_+^{k+1}(x_t) - (x_t^k - x_t^{k+1}) + (x_t^{k+1} - x_t^{k+2}) \\ &\geq \lambda_+^k(x_t) - (d+1)(x_t^k - x_t^{k+1}) + (x_t^{k+1} - x_t^{k+2}). \end{aligned}$$

Hence

$$\int_0^t \lambda_+^k(x_s) ds + \int_0^t (x_s^{k+1} - x_s^{k+2}) ds \leq e^{(d+1)t} (x_t^k - x_t^{k+1})$$

and

$$\int_0^t (x_s^k - x_s^{k+1}) ds \leq t e^{(d+1)t} (x_t^k - x_t^{k+1}).$$

Also

$$\int_0^t \lambda_+^k(x_s) ds \leq d \int_0^t (x_s^{k-1} - x_s^k) ds \leq d t e^{(d+1)t} (x_t^{k-1} - x_t^k).$$

Fix $\varepsilon > 0$ and set

$$f(t) = \sup_{k \in \mathbb{N}} \sup_{s \leq t} \mathbb{E}(|\gamma_s^k|^2) / \delta_s^k$$

where $\delta_t^k = \min\{x_t^{k-1} - x_t^k, x_t^k - x_t^{k+1}\} + \varepsilon$. Then $f(t_0) < \infty$. Note that

$$\begin{aligned} & \mathbb{E}(|\int_0^t \sigma_+^k(x_s) dB_+(s) - \int_0^t \sigma_-^k(x_s) dB_-(s)|^2) \\ &= \int_0^t \lambda_+^k(x_s) ds + \int_0^t (x_s^k - x_s^{k+1}) ds \leq (dt + 1) e^{(d+1)t} \delta_t^k. \end{aligned}$$

We have

$$\nabla b^k(x) \gamma = \lambda d (x^{k-1})^{d-1} \gamma^{k-1} - (\lambda d (x^k)^{d-1} + 1) \gamma^k + \gamma^{k+1}.$$

We will make use of the following estimates

$$\begin{aligned} & \mathbb{E}(|\int_0^t \lambda d (x_s^{k-1})^{d-1} \gamma_s^{k-1} ds|^2) \\ & \leq \int_0^t \lambda^2 d^2 (x_s^{k-1})^{2(d-1)} \delta_s^{k-1} ds \int_0^t \mathbb{E}(|\gamma_s^{k-1}|^2) / \delta_s^{k-1} ds \\ & \leq d^2 \int_0^t \lambda_+^k(x_s) ds \int_0^t f(s) ds \leq (d^2 + d^3 t) e^{(d+1)t} \delta_t^k \int_0^t f(s) ds \end{aligned}$$

and

$$\begin{aligned} & \mathbb{E}(|\int_0^t (\lambda d (x_s^k)^{d-1} + 1) \gamma_s^k ds|^2) \\ & \leq \int_0^t (\lambda d (x_s^k)^{d-1} + 1)^2 \delta_s^k ds \int_0^t \mathbb{E}(|\gamma_s^k|^2) / \delta_s^k ds \leq (d+1)^2 t e^{(d+1)t} \delta_t^k \int_0^t f(s) ds \end{aligned}$$

and

$$\mathbb{E}(|\int_0^t \gamma_s^{k+1} ds|^2) \leq \int_0^t \delta_s^{k+1} ds \int_0^t \mathbb{E}(|\gamma_s^{k+1}|^2) / \delta_s^{k+1} ds \leq e^{(d+1)t} \delta_t^k \int_0^t f(s) ds.$$

Now, from equation (13), for all $t \leq t_0$,

$$\mathbb{E}(|\gamma_t^k|^2) \leq A\delta_t^k + B\delta_t^k \int_0^t f(s) ds$$

where $A = 4(dt + 1)e^{(d+1)t}$ and $B = 8d(d + 1)(dt + 1)e^{(d+1)t}$. So $f(t) \leq Ae^{Bt}$ by Gronwall's lemma. This bound does not depend on ε , so the proposition follows by letting $\varepsilon \rightarrow 0$. $\square$

References

- [1] DARLING, R.W.R. AND NORRIS, J.R. (2002). Structure of large random hypergraphs. To appear in *Ann. Appl. Prob.*
- [2] DEIMLING, K. (1977). *Ordinary Differential Equations in Banach Spaces*. Lecture Notes in Mathematics 596, Springer-Verlag, Berlin.
- [3] EAGER, D.L., LAZOKWSKA, E.D. AND ZAHORJAN, J. (1986). Adaptive load sharing in homogeneous distributed systems. *IEEE Trans. Soft. Eng.* **12** 662–675.
- [4] ETHIER, S.N. AND KURTZ, T.G. (1986). *Markov Processes: Characterization and Convergence*. Wiley, New York.
- [5] GRAHAM, C. (2000). Kinetic limits for large communication networks. In *Modelling in Applied Sciences* (N. Bellomo and M. Pulvirenti, eds.) 317–370. Birkhäuser, Basel.
- [6] GRAHAM, C. (2000). Chaoticity on path space for a queueing network with selection of the shortest queue among several. *J. Appl. Prob.* **37** 198–201.
- [7] GRAHAM, C. (2004). Functional central limit theorems for a large network in which customers join the shortest of several queues. Preprint.

- [8] KOMLÓS, J., MAJOR, P. AND TUSNÁDY, G. (1975). An Approximation of Partial Sums of Independent RV's and the sample DF I. *Z. Wahrsch. verw. Gebiete* **32** 111–131.
- [9] LUCZAK, M.J. (2003). A quantitative law of large numbers via exponential martingales. In *Stochastic Inequalities and Applications* (E. Giné, C. Houdré and D. Nualart, eds.) Birkhäuser Series Progress in Probability **56** 93–111. Birkhäuser, Basel.
- [10] LUCZAK, M.J. AND MCDIARMID, C. (2003). On the power of two choices: balls and bins in continuous time. Preprint.
- [11] LUCZAK, M.J. AND MCDIARMID, C. (2004). On the maximum queue length in the supermarket model. Preprint.
- [12] MARTIN, J.B. AND SUHOV, Y.M. (1998). Fast Jackson networks. *Ann. Appl. Prob.* **9** 854–870.
- [13] MITZENMACHER, M. (1996). The power of two choices in randomized load balancing. PhD thesis, Berkeley.
- [14] REVUZ, D. AND YOR, M. (1991). *Continuous Martingales and Brownian Motion*. Springer-Verlag, Berlin.
- [15] VVEDENSKAYA, N.D., DOBRUSHIN, R.L. AND KARPELEVICH, F.I. (1996). Queueing system with selection of the shortest of two queues: an asymptotic approach. *Prob. Inf. Transm.* **32** 15–27.